

Anisotropic flow background to jet-correlation relative to reaction plane: a revisit of the mathematical framework

Joshua Konzer Fuqiang Wang

*Department of Physics, Purdue University, 525 Northwestern Ave., West
Lafayette, IN 47907, USA*

Abstract

We derive the flow-background formula for jet-correlation analysis with high p_T trigger particles in *any* azimuth window relative to reaction plane, extending the mathematic framework of previous study in [1].

Key words: heavy-ion; reaction plane; jet-correlation

PACS: 25.75.-q, 25.75.Dw

- 1 Strong interaction of hard-scattering partons with the high density medium
- 2 created in relativistic heavy-ion collisions results in modification to jet-correlation
- 3 with high transverse momentum (p_T) trigger particles due to jet energy loss [2,3,4].
- 4 The away-side correlation opposite to the trigger particle was found to be
- 5 strongly modified [5,6]; the correlation structure is significantly broadened,
- 6 and the lost energy is dissipated into low p_T particles [5] and appears to ex-

Email addresses: jrkonz@purdue.edu (Joshua Konzer), fqwang@purdue.edu (Fuqiang Wang).

7 cite Mach cone shock sound waves in the created medium [7]. The near-side
8 correlation was found to be extended to large pseudo-rapidity [5,8,9,10]. The
9 long-ranged correlation is dubbed as the “ridge”. The physics of the ridge is
10 not well understood.

11 In turn, jet-correlation measurement with high p_T trigger particles has become
12 a powerful tool to study medium properties [3,4]. Jet-correlation relative to
13 reaction plane provides further capability by exploiting the non-sphericity of
14 the initial nuclear overlap region [11]. It was found that the ridge magni-
15 tude drops significantly with trigger particles from in-plane to out-of-plane in
16 medium central Au+Au collisions [11]. In particular, with the trigger particle
17 pointing at an angle relative to the reaction plane in a non-central collision,
18 the medium thicknesses along the directions at the two sides of the away-side
19 parton are different. This asymmetry in the medium thickness should have
20 different effects on the gluons radiated off the away-side parton and can re-
21 sult in asymmetric correlation functions. A recent study [12] explored such
22 possibilities.

23 Recently Chiu and Hwa [13] proposed a physics mechanism for the observed
24 ridge as being a result of interactions between radiated gluons and medium
25 flow. They argue that when jet propagation is aligned with the medium flow
26 direction, small-angle radiated gluons are collimated with the jet direction,
27 while the radiated gluons are pushed away from the jet direction when jet
28 propagation is perpendicular to the medium flow direction [13]. This model of
29 gluon emission correlated with medium flow can explain the observed decrease
30 of ridge magnitude from in-plane to out-of-plane [11]. Furthermore, the model
31 predicts asymmetric azimuthal correlation functions for trigger particles in
32 quadrants I+III and quadrants II and IV, separately [13]. Such asymmetry

may not be restricted only to this particular model implementation, but a more general feature from interactions between jets and medium flow (e.g., see Refs. [14,15,16,17,18,19]). Clearly, experimental test of this physics mechanism is important to further our understanding of jet-medium interaction.

One important aspect of jet-correlation analysis is the subtraction of combinatorial background which itself has an azimuthal dependence due to anisotropic flow of background particles. The formulism of this flow background is given in Ref. [1] for trigger particles within a restricted azimuthal range relative to reaction plane, summed over all four quadrants. Due to reflection symmetry, flow backgrounds in quadrants I and III are identical and so are those in quadrants II and IV, but flow backgrounds in quadrants I+III and II+IV are not the same. The flow-background formula given in Ref. [1], summed over all four quadrants, cannot be readily used for jet-correlation analysis separating trigger particles in quadrants I+III and II+IV. In this note we derive the needed formula.

Due to event anisotropy, particle azimuthal distribution is given by

$$\frac{dN}{d\phi} = \frac{N}{2\pi} \left[1 + \sum_{k=1}^{\infty} 2v_k \cos k(\phi - \psi_{RP}) \right] \quad (1)$$

where ψ_{RP} is the reaction plane azimuthal angle, and v_k is the k th harmonic coefficient. Trigger-associated particle pair distribution is given by

$$\begin{aligned} \frac{d^4N}{d\psi_{RP}d\phi_t d\phi d\Delta\phi} &= \frac{1}{2\pi} \frac{N_t}{2\pi} \left[1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k(\phi_t - \psi_{RP}) \right] \times \\ &\quad \frac{N}{2\pi} \left[1 + \sum_{k=1}^{\infty} 2v_k \cos k(\phi - \psi_{RP}) \right] \delta(\Delta\phi - \phi + \phi_t) \quad (2) \end{aligned}$$

Integrating over trigger particle azimuthal angle ϕ_t within a slice $|\phi_t - (\psi_{RP} + \phi_s)| < c$, and associated particle ϕ and ψ_{RP} over full 2π , we have

$$\frac{dN}{d\Delta\phi} = \frac{N_t N}{(2\pi)^3} \int_0^{2\pi} d\psi_{RP} \int_{\psi_{RP}+\phi_s-c}^{\psi_{RP}+\phi_s+c} d\phi_t \left[1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k(\phi_t - \psi_{RP}) \right] \times \left[1 + \sum_{k=1}^{\infty} 2v_k \cos k(\Delta\phi + \phi_t - \psi_{RP}) \right] \quad (3)$$

54 However, we do not know the real reaction plane ψ_{RP} , but only the measured
 55 event plane ψ_{EP} , which is smeared from ψ_{RP} by probability function $\rho(\Delta\psi)$
 56 where $\Delta\psi = \psi_{EP} - \psi_{RP}$ and $\int_0^{2\pi} \rho(\Delta\psi) d\Delta\psi \equiv 1$. Experimentally trigger
 57 particles are selected within slice $|\phi_t - (\psi_{EP} + \phi_s)| < c$, hence

$$\frac{dN}{d\Delta\phi} = \frac{N_t N}{(2\pi)^3} \int_0^{2\pi} d\psi_{RP} \int_0^{2\pi} \rho(\Delta\psi) d\Delta\psi \int_{\psi_{RP}+\Delta\psi+\phi_s-c}^{\psi_{RP}+\Delta\psi+\phi_s+c} d\phi_t \times \left[1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k(\phi_t - \psi_{RP}) \right] \times \left[1 + \sum_{k=1}^{\infty} 2v_k \cos k(\Delta\phi + \phi_t - \psi_{RP}) \right]$$

58 The integrand is

$$1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k(\phi_t - \psi_{RP}) + \sum_{k=1}^{\infty} 2v_k \cos k(\Delta\phi + \phi_t - \psi_{RP}) + \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} 2v_i^{(t)} v_j \{ \cos [(j+i)(\phi_t - \psi_{RP}) + j\Delta\phi] + \cos [(j-i)(\phi_t - \psi_{RP}) + j\Delta\phi] \}$$

59 Integrating over ϕ_t , we obtain

$$\frac{dN}{d\Delta\phi} = \frac{2c N_t N}{(2\pi)^2} \int_0^{2\pi} \rho(\Delta\psi) d\Delta\psi \left\{ 1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \frac{\sin kc}{kc} \cos k(\phi_s + \Delta\psi) + \sum_{k=1}^{\infty} 2v_k \frac{\sin kc}{kc} \cos k(\Delta\phi + \phi_s + \Delta\psi) + \sum_{k=1}^{\infty} 2v_k^{(t)} v_k \cos k\Delta\phi + \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} 2v_i^{(t)} v_j \frac{\sin(j+i)c}{(j+i)c} \cos [(j+i)(\phi_s + \Delta\psi) + j\Delta\phi] + \sum_{j=1}^{\infty} \sum_{i=1, i \neq j}^{\infty} 2v_i^{(t)} v_j \frac{\sin(j-i)c}{(j-i)c} \cos [(j-i)(\phi_s + \Delta\psi) + j\Delta\phi] \right\} \quad (6)$$

60 Taking $\int_0^{2\pi} \rho(\Delta\psi) d\Delta\psi \cos n\Delta\phi \equiv \langle \cos n\Delta\phi \rangle$ and $\int_0^{2\pi} \rho(\Delta\psi) d\Delta\psi \sin n\Delta\phi = 0$,

61 we have

$$\begin{aligned}
\frac{dN}{d\Delta\phi} = \frac{2cN_tN}{(2\pi)^2} & \left\{ 1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k\phi_s \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \right. \\
& \sum_{k=1}^{\infty} 2v_k \cos k(\Delta\phi + \phi_s) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \sum_{k=1}^{\infty} 2v_k^{(t)} v_k \cos k\Delta\phi + \\
& \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} 2v_i^{(t)} v_j \cos [(j+i)\phi_s + j\Delta\phi] \frac{\sin(j+i)c}{(j+i)c} \langle \cos(j+i)\Delta\psi \rangle + \\
& \left. \sum_{j=1}^{\infty} \sum_{i=1, i \neq j}^{\infty} 2v_i^{(t)} v_j \cos [(j-i)\phi_s + j\Delta\phi] \frac{\sin(j-i)c}{(j-i)c} \langle \cos(j-i)\Delta\psi \rangle \right\}
\end{aligned}$$

62 The two $\sum \sum$ terms can be rewritten into:

$$\begin{aligned}
& \sum_{j=1}^{\infty} 2v_j \sum_{k=j+1}^{\infty} v_{k-j}^{(t)} \cos(k\phi_s + j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \\
& \sum_{j=1}^{\infty} 2v_j \sum_{k=1-j, k \neq 0}^{\infty} v_{k+j}^{(t)} \cos(k\phi_s - j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \\
& = \sum_{j=1}^{\infty} 2v_j \sum_{k=j+1}^{\infty} v_{k-j}^{(t)} \cos(k\phi_s + j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \\
& \sum_{j=1}^{\infty} 2v_j \left(\sum_{k=1}^{\infty} v_{k+j}^{(t)} + \sum_{k=1-j}^{-1} v_{k+j}^{(t)} \right) \cos(k\phi_s - j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \\
& = \sum_{j=1}^{\infty} 2v_j \left(\sum_{k=j+1}^{\infty} v_{k-j}^{(t)} + \sum_{k=1}^{j-1} v_{j-k}^{(t)} \right) \cos(k\phi_s + j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \\
& \sum_{j=1}^{\infty} 2v_j \sum_{k=1}^{\infty} v_{k+j}^{(t)} \cos(k\phi_s - j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \\
& = \sum_{j=1}^{\infty} 2v_j \sum_{k=1, k \neq j}^{\infty} v_{|k-j|}^{(t)} \cos(k\phi_s + j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \\
& \sum_{j=1}^{\infty} 2v_j \sum_{k=1}^{\infty} v_{k+j}^{(t)} \cos(k\phi_s - j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \tag{8}
\end{aligned}$$

63 Thus

$$\begin{aligned}
\frac{dN}{d\Delta\phi} = \frac{2cN_tN}{(2\pi)^2} & \left\{ 1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k\phi_s \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \right. \\
& \sum_{k=1}^{\infty} 2v_k \cos k(\Delta\phi + \phi_s) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \sum_{k=1}^{\infty} 2v_k^{(t)} v_k \cos k\Delta\phi + \\
& \sum_{j=1}^{\infty} 2v_j \sum_{k=1, k \neq j}^{\infty} v_{|k-j|}^{(t)} \cos(k\phi_s + j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle +
\end{aligned}$$

$$\sum_{j=1}^{\infty} 2v_j \sum_{k=1}^{\infty} v_{k+j}^{(t)} \cos(k\phi_s - j\Delta\phi) \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \Bigg\} \quad (9)$$

64 The 3rd and 5th term can be combined if using convention $v_0^{(t)} = 1$, yielding

$$\begin{aligned} \frac{dN}{d\Delta\phi} = \frac{2cN_t N}{(2\pi)^2} \Bigg\{ & 1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k\phi_s \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle + \sum_{k=1}^{\infty} 2v_k^{(t)} v_k \cos k\Delta\phi + \\ & \sum_{j=1}^{\infty} 2v_j \sum_{k=1}^{\infty} \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \left[v_{|k-j|}^{(t)} \cos(k\phi_s + j\Delta\phi) + v_{k+j}^{(t)} \cos(k\phi_s - j\Delta\phi) \right] \Bigg\} \end{aligned}$$

65 The number of trigger particles within the given slice is

$$\begin{aligned} N_t^{(R)} &= \frac{N_t}{(2\pi)^2} \int_0^{2\pi} d\psi \int_0^{2\pi} \rho(\Delta\psi) d\Delta\psi \int_{\psi_{RP} + \Delta\psi + \phi_s - c}^{\psi_{RP} + \Delta\psi + \phi_s + c} d\phi_t \left[1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k(\phi_t - \psi_{RP}) \right] \\ &= \frac{2cN_t}{(2\pi)^2} \int_0^{2\pi} d\psi \int_0^{2\pi} \rho(\Delta\psi) d\Delta\psi \left[1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \frac{\sin kc}{kc} \cos k(\Delta\psi + \phi_s) \right] \\ &= \frac{2cN_t}{2\pi} \left(1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k\phi_s \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \right) \end{aligned} \quad (11)$$

66 Normalized by the number of trigger particles, the correlation function be-

67 comes

$$\begin{aligned} \frac{1}{N_t^{(R)}} \frac{dN}{d\Delta\phi} &= \frac{N}{2\pi} \left(1 + \sum_{i=1}^{\infty} 2v_i \times \right. \\ & \quad \left. \frac{v_i^{(t)} \cos i\Delta\phi + \sum_{k=1}^{\infty} \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle \left[v_{|k-i|}^{(t)} \cos(i\Delta\phi + k\phi_s) + v_{k+i}^{(t)} \cos(i\Delta\phi - k\phi_s) \right]}{1 + \sum_{k=1}^{\infty} 2v_k^{(t)} \cos k\phi_s \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle} \right) \end{aligned} \quad (12)$$

68 Using shorthand notation

$$69 \quad R_k = \frac{\sin kc}{kc} \langle \cos k\Delta\psi \rangle, \quad (13)$$

70 we rewrite our result into

$$\frac{1}{N_t^{(R)}} \frac{dN}{d\Delta\phi} = \frac{N}{2\pi} \left(1 + \sum_{i=1}^{\infty} 2v_i \times \frac{v_i^{(t)} \cos i\Delta\phi + \sum_{k=1}^{\infty} R_k \left[v_{|k-i|}^{(t)} \cos(i\Delta\phi + k\phi_s) + v_{k+i}^{(t)} \cos(i\Delta\phi - k\phi_s) \right]}{1 + \sum_{k=1}^{\infty} 2v_k^{(t)} R_k \cos k\phi_s} \right) \quad (14)$$

At mid-rapidity, odd harmonics vanish. Thus the result is invariant under $\phi_s \rightarrow \phi_s + \pi$. The result is different under $\phi_s \rightarrow -\phi_s$; both $\frac{dN(\phi_s)}{d\Delta\phi}$ and $\frac{dN(-\phi_s)}{d\Delta\phi}$ are asymmetric about $\Delta\phi = 0$, but the average of the two,

$$\frac{1}{N_t^{(R)}} \frac{dN}{d\Delta\phi} = \frac{N}{2\pi} \left(1 + \sum_{i=\text{even}}^{\infty} 2v_i \frac{v_i^{(t)} + \sum_{k=\text{even}}^{\infty} (v_{|k-i|}^{(t)} + v_{k+i}^{(t)}) R_k \cos k\phi_s}{1 + \sum_{k=\text{even}}^{\infty} 2v_k^{(t)} R_k \cos k\phi_s} \cos i\Delta\phi \right) \quad (15)$$

is symmetric. Eq. 15 recovers the result in Ref. [1].

Keeping terms up to $v_2 v_4$ and v_6 , Eq. 14 becomes

$$\frac{1}{N_t^{(R)}} \frac{dN}{d\Delta\phi} = \frac{N}{2\pi} \left\{ 1 + \frac{1}{1 + \sum_{k=2,4,6} 2v_k^{(t)} R_k \cos k\phi_s} \times \left(\begin{aligned} &2v_2 R_2 \cos 2(\Delta\phi + \phi_s) + 2v_4 R_4 \cos 4(\Delta\phi + \phi_s) + \\ &2v_6 R_6 \cos 6(\Delta\phi + \phi_s) + 2v_2 v_2^{(t)} [\cos 2\Delta\phi + R_4 \cos 2(\Delta\phi + 2\phi_s)] \\ &+ 2v_2 v_4^{(t)} [R_2 \cos 2(\Delta\phi - \phi_s) + R_6 \cos 2(\Delta\phi + 2\phi_s)] + \\ &2v_2^{(t)} v_4 [R_2 \cos 2(2\Delta\phi + \phi_s) + R_6 \cos 2(2\Delta\phi + 3\phi_s)] \end{aligned} \right) \right\} \quad (16)$$

For illustration we calculate the flow modulation, the quantity in the curly brackets in Eq. (16), for typical elliptic flow magnitudes of $v_2^{(t)} = 0.10$ and $v_2 = 0.05$, and $v_4^{(t)} = (v_2^{(t)})^2$, $v_6^{(t)} = (v_2^{(t)})^3$, $v_4 = v_2^2$ and $v_6 = v_2^3$. We shall take the

80 event plane resolutions to be $\langle \cos 2\Delta\psi \rangle = 1$, $\langle \cos 4\Delta\psi \rangle = 1$, and $\langle \cos 6\Delta\psi \rangle =$
 81 1 (i.e., perfect event plane reconstruction). For smaller resolutions, the flow
 82 modulation is reduced. The calculated flow modulation is shown in Fig. 1(a)
 83 in the solid curve for trigger particle azimuthal angle range $15^\circ < \phi_t < 30^\circ$
 84 (i.e., $\phi_s = 22.5^\circ, c = 7.5^\circ$). Also shown are the individual contributions of the
 85 order v_2 , v_4 , and v_6 , in dashed, dotted, and dash-dotted curves, respectively.
 86 In Fig. 1(b) we show the flow modulations for ranges $30^\circ < \phi_t < 45^\circ$ (dashed
 87 curve), $-45^\circ < \phi_t < -30^\circ$ (dotted curve), and the combined range $30^\circ <$
 88 $|\phi_t| < 45^\circ$ (solid curve).




Fig. 1. (a) Flow modulation, the quantity in the curly brackets in Eq. (16)
 for trigger particle azimuthal angle range $15^\circ < \phi_t < 30^\circ$ (solid curve).
 Typical elliptic flow magnitudes of $v_2^{(t)} = 0.10$ and $v_2 = 0.05$, and
 $v_4^{(t)} = (v_2^{(t)})^2, v_6^{(t)} = (v_2^{(t)})^3, v_4 = v_2^2$ and $v_6 = v_2^3$ are used and event plane res-
 olutions are taken to be $\langle \cos 2\Delta\psi \rangle = \langle \cos 4\Delta\psi \rangle = \langle \cos 6\Delta\psi \rangle = 1$. The dashed,
 dotted, and dash-dotted curves are the individual contributions of the order v_2 ,
 v_4 , and v_6 , respectively. (b) Flow modulations for ranges $30^\circ < \phi_t < 45^\circ$ (dashed
 curve), $-45^\circ < \phi_t < -30^\circ$ (dotted curve), and the combined range $30^\circ < |\phi_t| < 45^\circ$
 (solid curve).

89 In summary, we have derived the flow-background formula for jet-correlation
 90 analysis with high p_T trigger particles in *any* azimuth window relative to
 91 reaction plane, extending the mathematic framework of previous study in [1].

92 Our main result is in Eq. (14). An approximation up to the order of v_2v_4 (and
93 v_6) is given in Eq. (16).

94 *Acknowledgment*– This work is supported by U.S. Department of Energy under
95 Grant DE-FG02-88ER40412.

96 References

- 97 [1] J. Bielcikova *et al.*, Phys. Rev. C **69**, 021901(R) (2004).
- 98 [2] X.-N. Wang and M. Gyulassy, Phys. Rev. Lett. **68** (1992) 1480.
- 99 [3] J. Adams *et al.* (STAR Collaboration), Nucl. Phys. **A757** (2005) 102.
- 100 [4] K. Adcox *et al.* (PHENIX Collaboration) Nucl. Phys. **A757** (2005) 184.
- 101 [5] J. Adams *et al.* (STAR Collaboration), Phys. Rev. Lett. **95** (2005) 152301.
- 102 [6] A. Adare *et al.* (PHENIX Collaboration), Phys. Rev. C **78** (2008) 014901.
- 103 [7] B. I. Abelev *et al.* (STAR Collaboration), Phys. Rev. Lett. **102** (2009) 052302.
- 104 [8] J. Putschke (STAR Collaboration), J. Phys. G **34** (2007) S679.
- 105 [9] L. Molnar (STAR Collaboration), J. Phys. G **34** (2007) S593.
- 106 [10] B. Alver *et al.*, arXiv:0903.2811.
- 107 [11] A. Feng (STAR Collaboration), J. Phys. G **35** (2008) 104082.
- 108 [12] J. Jia, S. Esumi, and R. Wei, arXiv:0903.3263.
- 109 [13] C. B. Chiu and R. C. Hwa, arXiv:0809.3018.
- 110 [14] S. A. Voloshin, Phys. Lett. B **632** (2006) 490.
- 111 [15] E. V. Shuryak, Phys. Rev. C **76** (2007) 047901.

- 112 [16] A. Dumitru, F. Gelis, L. McLerran, and R. Venugopalan, Nucl. Phys. **A810**
113 (2008) 91.
- 114 [17] S. Gavin, L. McLerran, and G. Moschelli, arXiv:0806.4718.
- 115 [18] S. J. Lindenbaum and R. S. Longacre, Phys. Rev. C **78** (2008) 054904.
- 116 [19] J. Takahashi *et al.*, arXiv:0902.4870.